
In situ plankton imaging with a UVP (Underwater Vision Profiler)




UVP - Background

The UVP (Underwater Vision Profiler) was developed by Gaby Gorsky's lab in Villefranchesur-mer, France (Picheral et al. 2010) and has evolved to the current generation, UVP6 (Picheral et al. 2022). In the CCE-LTER *California Current Ecosystem* Long-Term Ecological Research site, we have been using the UVP5 on our process cruises since 2008. Here we present a few of the results from our use of that instrument, thanks largely to Tristan Biard, a former SIO postdoc who has worked extensively with UVP images and data.


CCE-LTER has recently purchased a UVP6-HF (highfrequency sampling) for deployment on a CTDrosette, thus permitting a vertical profile of images of zooplankton and larger suspended particles to be made each time the CTD is lowered.





Siphonophore





Biard T, Krause JW, Stukel MR, Ohman MD (2018) The Significance of Giant Phaeodarians (Rhizaria) to Biogenic Silica Export in the California Current Ecosystem. Global Biogeochemical Cycles 32: 987-1004 doi 10.1029/2018gb005877

Biard T, Ohman MD (2020) Vertical niche definition of test-bearing protists (Rhizaria) into the twilight zone revealed by in situ imaging. Limnology and Oceanography doi 10.1002/lno.11472 Biard T, Stemmann L, Picheral M, Mayot N, Vandromme P, Hauss H, Gorsky G, Guidi L, Kiko R, Not F (2016) In situ imaging reveals the biomass of giant protists in the global ocean. Nature 532: 504-+ doi 10.1038/nature17652 Ellen JS, C. A. Graaf, and M. D. Ohman (2019) Improving plankton image classification using context metadata. Limnology and Oceanography - Methods 17: 439-461 doi 10.1002/lom3.10324 Ellen J, Ohman MD (2022) Beyond Transfer Learning: Leveraging external images to improve plankton image classification. 2022 Ocean Sciences Meeting, Honolulu, HI

Karstensen J, Kim D-g, Lekanoff RM, Lombard F, Lopes RM, Marec C, McDonnell AMP, Niemeyer D, Noyon M, O'Daly SH, Ohman MD, Pretty JL, Rogge A, Searson S, Shibata M, Tanaka Y, Tanhua T, Taucher J, Trudnowska E, Turner JS, Waite AM, Stemmann L (2022) A global marine particle size distribution dataset obtained with the Underwater Vision Profiler 5. Earth System Science Data 14: 1-22 doi https://doi.org/10.5194/essd-14-1-2022 L (2022) The Underwater Vision Profiler 6: an imaging sensor of particle size spectra and plankton, for autonomous and cabled platforms. Limnology and Oceanography-Methods 20: 115-129 doi 10.1002/lom3.10475

Kiko R, Picheral M, Antoine D, Babin M, Berline L, Biard T, Boss E, Brandt P, Carlotti F, Christiansen S, Coppola L, de la Cruz L, Diamond-Riquier E, de Madron XD, Elineau A, Gorsky G, Guidi L, Hauss H, Irisson J-O, Karp-Boss L, Picheral M, Catalano C, Brousseau D, Claustre H, Coppola L, Leymarie E, Coindat JM, Dias F, Fevre S, Guidi L, Irisson JO, Legendre L, Lombard F, Mortier L, Penkerch C, Rogge A, Schmechtig C, Thibault S, Tixier T, Waite A, Stemmann Picheral M, Guidi L, Stemmann L, Karl DM, Iddaoud G, Gorsky G (2010) The Underwater Vision Profiler 5: An advanced instrument for high spatial resolution studies of particle size spectra and zooplankton. Limnology and *Oceanography-Methods* 8: 462-473 doi 10.4319/lom.2010.8.462


Stukel MR, Biard T, Krause J, Ohman MD (2018) Large Phaeodaria in the twilight zone: Their role in the carbon cycle. Limnology and Oceanography 63: 2579-2594 doi 10.1002/lno.10961

Mark D. Ohman¹, Moira Décima¹, Tristan Biard²

¹Scripps Institution of Oceanography, UCSD ² Laboratoire d'Océanologie et de Géosciences, University of Lille, Wimereux, France

CCE-LTER seeks to collaborate with CalCOFI to make regular deployments of the UVP6-HF on CalCOFI cruises. No additional wire time or ship time is required for such deployments, which will enable in situ imaging of zooplankton and detrital material with each CTD cast. This addition would make it possible to resolve vertical structure of planktonic organisms critical to the food web and to biogeochemical cycles. These analyses would be complementary to the vertically integrated plankton tows (210-0 m), for which only a small subset of the samples can currently be analyzed. The UVP6-HF can resolve suspended marine particles as small as 0.1 mm, size spectra of the plankton community, and provides a better assessment of fragile, gelatinous zooplankton than conventional net samples.

Advanced Machine Learning tools (e.g., Ellen et al. 2019, 2022) are now available to facilitate the rapid classification of plankton images.



References

The Opportunity

Representative Images and Results


We express thanks to the National Science Foundation for support of the CCE-LTER site and for the purchase of a UVP6-HF.

We also thank Marc Picheral and the LOV (Laboratoire d'Océanographie de Villefranche), France, as well as Lee Karp-Boss and Emmanuel Boss of the University of Maine, for loaning us UVP instruments.






Rhizaria (Protists)



Vertical niches of planktonic Rhizaria

Vertical profiles of Aulosphaeridae (Phaeodaria)



Acknowledgments

