

Part IV

## INDEX TO CALCOFI REPORTS 1983–1988

This index is in two parts: author-title and subject. The index was generated from the publications listed on the right. The year in parentheses is the publication date. The volume number (roman numerals) is used to identify the references in the index entries. An index to previous issues, 1950–1982, appeared in volume XXIV.

### AUTHOR-TITLE INDEX

- Ahlstrom, E. H.: Descriptions of larvae of California yellowtail, *Seriola lalandei*, and three other carangids from the eastern tropical Pacific: *Chloroscombrus orqueta*, *Caranx caballus*, and *Caranx sexfasciatus*, XXVI:139–159
- Alarcón, V. H.: Spawning frequency and sex ratio in the Peruvian anchovy, *Engraulis ringens*, XXV:43–52
- Alheit, J., V. H. Alarcón, and B. J. Macewicz: Spawning frequency and sex ratio in the Peruvian anchovy, *Engraulis ringens*, XXV:43–52
- Allen, L. G.: Diel variation in catch parameters for fishes sampled by 7.6-m otter trawl in southern California coastal waters, XXV:119–134
- Alvarez, L. G., A. Badan-Dangon, and J. M. Robles: Lagrangian observations of near-surface currents in Canal de Ballenas, XXV:35–42
- Ambrose, D. A.: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182–193
- : Development and distribution of larvae and pelagic juveniles of ocean whitefish, *Caulolatilus princeps*, in the CalCOFI survey region, XXVII:162–169
- Ambrose, D. A., J. L. Butler, H. G. Moser, B. Y. Sumida, E. M. Sandknop, and E. G. Stevens: Description of the larvae of the cusk eels *Ophidion scrippsae* and *Chilara taylori* (family Ophidiidae), XXIV:226–234
- Anderson, S. L., L. W. Botsford, and W. H. Clark, Jr.: Size distributions and sex ratios of ridgeback prawns, *Sicyonia ingentis*, in the Santa Barbara Channel (1979–1981), XXVI:169–174
- Antonelis, G. A., Jr. and M. A. Perez: Estimated annual food consumption by northern fur seals in the California Current, XXV:135–145
- Arcos, F., and A. Fleminger: Distribution of filter-feeding calanoid copepods in the eastern equatorial Pacific, XXVII:170–187
- Armstrong, M. J.: The influence of large-scale environmental processes on neritic fish populations in the Benguela Current system, XXVI:72–92
- Armstrong, M., P. Shelton, I. Hampton, G. Jolly, and Y. Melo: Egg production estimates of anchovy biomass in the southern Benguela system, XXIX:137–157
- Arvizu-Martinez, J.: Fisheries activities in the Gulf of California, Mexico, XXVIII:32–36
- Badan-Dangon, A.: Coupling of the Pacific sardine, *Sardinops sagax caeruleus*, life cycle with the Gulf of California pelagic environment, XXIX:102–109
- California Cooperative Oceanic Fisheries Investigations Reports:  
Volume XXIV, January 1 to December 31, 1982 (1983)  
Volume XXV, January 1 to December 31, 1983 (1984)  
Volume XXVI, January 1 to December 31, 1984 (1985)  
Volume XXVII, January 1 to December 31, 1985 (1986)  
Volume XXVIII, January 1 to December 31, 1986 (1987)  
Volume XXIX, January 1 to December 31, 1987 (1988)
- : Lagrangian observations of near-surface currents in Canal de Ballenas, XXV:35–42
- Bagarinao, T., and J. R. Hunter: The visual feeding threshold and action spectrum of northern anchovy, *Engraulis mordax*, larvae, XXIV:245–254
- Bailey, K. M., R. C. Francis, and K. F. Mais: Evaluating incidental catches of 0-age Pacific hake to forecast recruitment, XXVII:109–112
- Bakun, A.: Comparative studies and the recruitment problem: searching for generalizations, XXVI:30–40
- Barber, R. T., J. E. Kogelschatz, and F. P. Chavez: Origin of productivity anomalies during the 1982–83 El Niño, XXVI:65–71
- Baumgartner, T. R.: Coupling of the Pacific sardine, *Sardinops sagax caeruleus*, life cycle with the Gulf of California pelagic environment, XXIX:102–109
- Bedford, D. W.: The biology of three pelagic sharks from California waters, and their emerging fisheries: a review, XXIV:57–69
- Bedford, D. W., and F. B. Hagerman: The billfish fishery resource of the California Current, XXIV:70–78
- Bernard, H. J., J. B. Hedgepeth, and S. B. Reilly: Stomach contents of albacore, skipjack, and bonito caught off southern California during summer 1983, XXVI:175–182
- Bindman, A. G.: The 1985 spawning biomass of the northern anchovy, XXVII:16–24
- Botsford, L. W.: Size distributions and sex ratios of ridgeback prawns, *Sicyonia ingentis*, in the Santa Barbara Channel (1979–1981), XXVI:169–174
- Boyd, A. J.: The influence of large-scale environmental processes on neritic fish populations in the Benguela Current system, XXVI:72–92
- Brainard, R. E.: Anomalous warm events in eastern boundary current systems, XXVI:51–64
- Brewer, G. D.: Nearshore production of young anchovy, XXIV:235–244
- Brinton, E., A. Fleminger, and D. Siegel-Causey: The temperate and tropical planktonic biotas of the Gulf of California, XXVII:228–266
- Brooks, E. R., and M. M. Mullin: Diel changes in the vertical distribution of biomass and species in the Southern California Bight, XXIV:210–215
- Busby, M. S.: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182–193

- Butler, J. L.: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182–193
- : Description of the larvae of the cusk eels *Ophiodon scrippsae* and *Chilara taylora* (family Ophidiidae), XXIV:226–234
- Butler, J. L., and B. Rojas de Mendiola: Growth of larval sardines off Peru, XXVI:113–118
- Cailliet, G. M., and D. W. Bedford: The biology of three pelagic sharks from California waters, and their emerging fisheries: a review, XXIV:57–69
- Chavez, F. P.: Origin of productivity anomalies during the 1982–83 El Niño, XXVI:65–71
- Chelton, D. B.: Zooplankton variability in the California Current, 1951–1982, XXVIII:59–96
- Chen, Y.: The vertical distribution of some pelagic copepods in the eastern tropical Pacific, XXVII:205–227
- Chess, J. R., S. E. Smith, and P. C. Fischer: Trophic relationships of the shortbelly rockfish, *Sebastes jordani*, off central California, XXIX:129–136
- Clark, W. H., Jr.: Size distributions and sex ratios of ridgeback prawns, *Sicyonia ingentis*, in the Santa Barbara Channel (1979–1981), XXVI:169–174
- Crooke, S. J.: Yellowtail, *Seriola lalandei* Valenciennes, XXIV:84–87
- Cross, J. N.: Demersal fishes of the upper continental slope off southern California, XXVIII:155–167
- DeMartini, E. E.: The feeding habits and distribution of juvenile–small adult California halibut, *Paralichthys californicus*, in coastal waters off northern San Diego County, XXIV:194–201
- : The feeding habits of juvenile–small adult barred sand bass, *Paralabrax nebulifer*, in nearshore waters off northern San Diego County, XXV:105–111
- : Tests of ovary subsampling options and preliminary estimates of batch fecundity for two Paralabrax species, XXVIII:168–170
- DeMartini, E. E., and L. G. Allen: Diel variation in catch parameters for fishes sampled by a 7.6-m otter trawl in southern California coastal waters, XXV:119–134
- Deweese, M. C., and E. M. Strange: Drift bottle observations of the nearshore surface circulation off California, 1977–1983, XXV:68–73
- Dryden, K. A.: A historical review of fisheries statistical and environmental and societal influences off the Palos Verdes Peninsula, California, XXVIII:135–154
- Dugdale, R. C.: The effects of varying nutrient concentrations on biological production in upwelling regions, XXVI:93–96
- Dykstra, D. H., R. S. Grove, and J. B. Palmer: Coastal front observations with an infrared scanner, XXV:59–67
- Eber, L. E.: Larval fish assemblages in the California Current region, 1954–1960, a period of dynamic environmental change, XXVIII:97–127
- Espino, M. A., and C. Wosnitza-Mendo: Peruvian hake fisheries from 1971 to 1982, XXVII:113–120
- Fiedler, P. C.: Offshore entrainment of anchovy spawning habitat, eggs, and larvae by a displaced eddy in 1985, XXVII:144–152
- : Satellite remote sensing of the habitat of spawning anchovy in the Southern California Bight, XXIV:202–209
- : Some effects of El Niño 1983 on the northern anchovy, XXV:53–58
- Fischer, P. C.: Trophic relationships of the shortbelly rockfish, *Sebastes jordani*, off central California, XXIX:129–136
- Fleminger, A.: Distribution of filter-feeding calanoid copepods in the eastern equatorial Pacific, XXVII:170–187
- : The temperate and tropical planktonic biotas of the Gulf of California, XXVII:228–266
- Francis, R. C.: Evaluating incidental catches of 0-age Pacific hake to forecast recruitment, XXVII:109–112
- Gomez-Muñoz, V. M.: Modelo polinomial para crecimiento individual de larvas de la anchoveta norteña, *Engraulis mordax*, XXVII:141–143
- Gregory, P. A.: A historical review of fisheries statistics and environmental and societal influences off the Palos Verdes Peninsula, California, XXVIII:135–154
- Grove, R. S.: Coastal front observations with an infrared scanner, XXV:59–67
- : Physical-chemical characteristics and zooplankton biomass on the continental shelf off southern California, XXVII:37–52
- Hagerman, F. B.: The billfish fishery resource of the California Current, XXIV:70–78
- Hammann, M. G., T. R. Baumgartner, and A. Badan-Dangon: Coupling of the Pacific sardine, *Sardinops sagax caeruleus*, life cycle with the Gulf of California pelagic environment, XXIX:102–109
- Hampton, I.: Egg production estimates of anchovy biomass in the southern Benguela system, XXIX:137–157
- Hanan, D. A.: Bluefin tuna in the eastern North Pacific Ocean, XXIV:97–98
- Hayward, T. L.: Determining chlorophyll on the 1984 CalCOFI surveys, XXV:74–79
- : Evaluation of some techniques for preserving nutrients in stored seawater samples, XXVI:160–168
- Hedgecock, D., K. Nelson, and L. G. Lopez-Lemus: Biochemical genetic and morphological divergence among three species of thread herring, *Opisthonema*, in northwest Mexico, XXIX:110–121
- Hedgepeth, J. B.: Stomach contents of albacore, skipjack, and bonito caught off southern California during summer 1983, XXVI:175–182
- Hernández-Vasquez, S.: Pesquerías pelágicas y neríticas de la costa occidental de Baja California, México, XXVIII:53–56
- Hewitt, R. P.: Anchovy egg dispersal and mortality as inferred from close-interval observations, XXVI:97–110
- : Historical review of the oceanographic approach to fishery research, XXIX:27–41
- : The 1984 spawning biomass of the northern anchovy, XXVI:17–25
- : The northern anchovy spawning biomass for the 1982–83 California fishing season, XXIV:16–27
- Hewitt, R. P., and G. D. Brewer: Nearshore production of young anchovy, XXIV:235–244
- Huato-Soberanis, L., and D. Lluch-Belda: Mesoscale cycles in the series of environmental indices related to the sardine fishery in the Gulf of California, XXVIII:128–134
- Hunter, J. R.: The visual feeding threshold and action spectrum of northern anchovy, *Engraulis mordax*, larvae, XXIV:245–254
- Jahn, A. E.: Abundance of southern California nearshore ichthyoplankton: 1978–1984, XXVII:53–64
- : Physical-chemical characteristics and zooplankton biomass on the continental shelf off southern California, XXVII:37–52
- : Sampling for eggs of sardine and other fishes in the coastal zone using the CalVET net, XXVIII:178–182
- Jahn, A. E., and P. E. Smith: Effects of sample size and contagion on estimating fish egg abundance, XXVIII:171–177
- Jimenez-Perez, L. C., and J. R. Lara-Lara: Zooplankton bio-

- mass and copepod community structure in the Gulf of California during the 1982–1983 El Niño event, XXIX:122–128
- Jolly, G.: Egg production estimates of anchovy biomass in the southern Benguela system, XXIX:137–157
- Jordan, G. A.: Can we relate larval fish abundance to recruitment or population stability? A preliminary analysis of recruitment to a temperate rocky reef, XXVII:65–83
- Kimball, A. S.: Comparative quality of rotifers and copepods as foods for larval fishes, XXV:81–86
- Klingbeil, R. A.: Pacific mackerel: a resurgent resource and fishery of the California Current, XXIV:35–45  
———: Recent increased abundance and potential productivity of Pacific mackerel, *Scomber japonicus*, XXVI:119–129
- Knight, M. D.: Variation in larval morphogenesis within the Southern California Bight population of *Euphausia pacifica* from winter through summer, 1977–1978, XXV:87–99
- Kogelschatz, J. E.: Origin of productivity anomalies during the 1982–83 El Niño, XXVI:65–71
- Lara-Lara, J. R.: Zooplankton biomass and copepod community structure in the Gulf of California during the 1982–1983 El Niño event, XXIX:122–128
- Laur, R. M.: The North Pacific albacore—an important visitor to California Current waters, XXIV:99–106
- Lavenberg, R. J.: Physical-chemical characteristics and zooplankton biomass on the continental shelf off southern California, XXVII:37–52
- Lavenberg, R. J., A. E. Jahn, G. E. McGowen, and J. H. Petersen: Sampling for eggs of sardine and other fishes in the coastal zone using the CalVET net, XXVIII:178–182
- Lavenberg, R. J., G. E. McGowen, A. E. Jahn, J. H. Petersen, and T. C. Sciarrotta: Abundance of southern California nearshore ichthyoplankton: 1978–1984, XXVII:53–64
- Leong, R.: Growth of jack mackerel, *Trachurus symmetricus*, in captivity, XXV:146–150
- Lluch-Belda, D.: Mesoscale cycles in the series of environmental indices related to the sardine fishery in the Gulf of California, XXVIII:128–134
- Lluch-Belda, D., F. J. Magallon, and R. A. Schwartzlose: Large fluctuations in the sardine fishery in the Gulf of California: possible causes, XXVII:136–140
- Loeb, V. J., P. E. Smith, and H. G. Moser: Geographical and seasonal patterns of larval fish species structure in the California Current area, 1975, XXIV:132–151  
———: Ichthyoplankton and zooplankton abundance patterns in the California Current area, 1975, XXIV:109–131  
———: Recurrent groups of larval fish species in the California Current area, XXIV:152–164
- Lopez-Lemus, L. G.: Biochemical genetic and morphological divergence among three species of thread herring, *Opisthonema*, in northwest Mexico, XXIX:110–121
- Love, M. S., J. S. Stephens, Jr., P. A. Morris, M. M. Singer, M. Sandhu, and T. C. Sciarrotta: Inshore soft substrata fishes in the Southern California Bight: an overview, XXVII:84–106
- MacCall, A. D.: Revised estimates of historical spawning biomass of the Pacific mackerel, *Scomber japonicus*, XXIX:81–90
- MacCall, A. D., R. A. Klingbeil, and R. D. Methot: Recent increased abundance and potential productivity of Pacific mackerel, *Scomber japonicus*, XXVI:119–129
- MacCall, A. D., and M. H. Prager: Historical changes in abundance of six fish species off southern California, based on CalCOFI egg and larva samples, XXIX:91–101
- MacCall, A. D., and G. D. Stauffer: Biology and fishery potential of jack mackerel, *Trachurus symmetricus*, XXIV:46–56
- Macewicz, B. J.: Spawning frequency and sex ratio in the Peruvian anchovy, *Engraulis ringens*, XXV:43–52
- MacGregor, J. S.: Growth of the blue rockfish, *Sebastes mystinus*, XXIV:216–225  
———: Relative abundance of four species of *Sebastes* off California and Baja California, XXVII:121–135
- Magallon, F. J.: Large fluctuations in the sardine fishery in the Gulf of California: possible causes, XXVII:136–140
- Magallón-Barajas, F. J.: The Pacific shrimp fishery of Mexico, XXVIII:43–52
- Mais, K. F.: Evaluating incidental catches of 0-age Pacific hake to forecast recruitment, XXVII:109–112
- McGowen, G. E.: Abundance of southern California nearshore ichthyoplankton: 1978–1984, XXVII:53–64  
———: Can we relate larval fish abundance to recruitment or population stability? A preliminary analysis of recruitment to a temperate rocky reef, XXVII:65–83  
———: Physical-chemical characteristics and zooplankton biomass on the continental shelf off southern California, XXVII:37–52  
———: Sampling for eggs of sardine and other fishes in the coastal zone using the CalVET net, XXVIII:178–182
- McLain, D. R., R. E. Brainard, and J. G. Norton: Anomalous warm events in eastern boundary current systems, XXVI:51–64
- McLain, D. R., and D. H. Thomas: Year-to-year fluctuations of the California Countercurrent and effects on marine organisms, XXIV:165–181
- Melo, Y.: Egg production estimates of anchovy biomass in the southern Benguela system, XXIX:137–157
- Methot, R. D.: Frame trawl for sampling pelagic juvenile fish, XXVII:267–278  
———: Recent increased abundance and potential productivity of Pacific mackerel, *Scomber japonicus*, XXVI:119–129
- Morris, P. A.: Can we relate larval fish abundance to recruitment or population stability? A preliminary analysis of recruitment to a temperate rocky reef, XXVII:65–83  
———: Inshore soft substrata fishes in the Southern California Bight: an overview, XXVII:84–106
- Moser, H. G.: CalCOFI time series: an overview of fishes, XXIX:66–78  
———: Description of the larvae of the cusk eels *Ophidion scrippsae* and *Chilara taylori* (family Ophidiidae), XXIV:226–234  
———: Descriptions of larvae of California yellowtail, *Seriola lalandei*, and three other carangids from the eastern tropical Pacific: *Chloroscombrus orqueta*, *Caranx caballus*, and *Caranx sexfasciatus*, XXVI:139–159  
———: Food and feeding of bocaccio, *Sebastes paucispinis*, and comparison with Pacific hake, *Merluccius productus*, larvae in the California Current, XXV:112–118  
———: Geographical and seasonal patterns of larval fish species structure in the California Current area, 1975, XXIV:132–151  
———: Ichthyoplankton and zooplankton abundance patterns in the California Current area, 1975, XXIV:109–131  
———: Recurrent groups of larval fish species in the California Current area, XXIV:152–164
- Moser, H. G., D. A. Ambrose, M. S. Busby, J. L. Butler, E. M. Sandknop, B. Y. Sumida, and E. G. Stevens: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182–193
- Moser, H. G., P. E. Smith, and L. E. Eber: Larval fish assemblages in the California Current region, 1954–1960, a period of dynamic environmental change, XXVIII:97–127

- Moser, H. G., B. Y. Sumida, D. A. Ambrose, E. M. Sandknop, and E. G. Stevens: Development and distribution of larvae and pelagic juveniles of ocean whitefish, *Caulolatilus princeps*, in the CalCOFI survey region, XXVII:162–169
- Muhlia-Melo, A.: The Mexican tuna fishery, XXVIII:37–42
- Mullin, M. M.: Diel changes in the vertical distribution of biomass and species in the Southern California Bight, XXIV:210–215
- Nelson, K.: Biochemical genetic and morphological divergence among three species of thread herring, *Opisthonema*, in northwest Mexico, XXIX:110–121
- Norton, J. G.: Anomalous warm events in eastern boundary current systems, XXVI:51–64
- Palmer, J. B.: Coastal front observations with an infrared scanner, XXV:59–67
- Perez, M. A.: Estimated annual food consumption by northern fur seals in the California Current, XXV:135–145
- Petersen, J. H.: Abundance of southern California nearshore ichthyoplankton: 1978–1984, XXVII:53–64
- : Sampling for eggs of sardine and other fishes in the coastal zone using the CalVET net, XXVIII:178–182
- Petersen, J. H., A. E. Jahn, R. J. Lavenberg, G. E. McGowen, and R. S. Grove: Physical-chemical characteristics and zooplankton biomass on the continental shelf off southern California, XXVII:37–52
- Phillips, G. M.: Interactions between fishing effort and catch per unit effort in the San Pedro fishery: a statistical note, XXIV:255–258
- Picaut, J.: Major dynamics affecting the eastern tropical Atlantic and Pacific oceans, XXVI:41–50
- Picquelle, S. J., and R. P. Hewitt: The 1983 spawning biomass for the northern anchovy, XXV:16–27
- The northern anchovy spawning biomass for the 1982–83 California fishing season, XXIV:16–27
- Plummer, K. M.: The feeding habits of juvenile-small adult barred sand bass, *Paralabrax nebulifer*, in nearshore waters off northern San Diego County, XXV:105–111
- Plummer, K. M., E. E. DeMartini, and D. A. Roberts: The feeding habits and distribution of juvenile-small adult California halibut, *Paralichthys californicus*, in coastal waters off northern San Diego County, XXIV:194–201
- Prager, M. H.: Historical changes in abundance of six fish species off southern California, based on CalCOFI egg and larva samples, XXIX:91–101
- Prager, M. H., and A. D. MacCall: Revised estimates of historical spawning biomass of the Pacific mackerel, *Scomber japonicus*, XXIX:81–90
- Quiñonez-Velázquez, C., and V. M. Gomez-Muñoz: Modelo polinomial para crecimiento individual de larvas de la anchoveta norteña, *Engraulis mordax*, XXVII:141–143
- Reed, R. J.: White seabass, *Atractoscion nobilis*, in California-Mexican waters: status of the fishery, XXIV:79–83
- Reed, R. J., and A. D. MacCall: Changing the size limit: how it could affect California halibut fisheries, XXIX:158–166
- Reid, J. L.: Physical oceanography, 1947–1987, XXIX:42–65
- Reilly, S. R.: Stomach contents of albacore, skipjack, and bonito caught off southern California during summer 1983, XXVI:175–182
- Roberts, D. A.: The feeding habits and distribution of juvenile-small adult California halibut, *Paralichthys californicus*, in coastal waters off northern San Diego County, XXIV:194–201
- Roberts, D. A., E. E. DeMartini, and K. M. Plummer: The feeding habits of juvenile-small adult barred sand bass, *Paralabrax nebulifer*, in nearshore waters off northern San Diego County, XXV:105–111
- Robles, J. M.: Lagrangian observations of near-surface currents in Canal de Ballenas, XXV:35–42
- Roesler, C. S., and D. B. Chelton: Zooplankton variability in the California Current, 1951–1982, XXVIII:59–96
- Rojas de Mendiola, B.: Growth of larval sardines off Peru, XXVI:113–118
- Sandhu, M.: Inshore soft substrata fishes in the Southern California Bight: an overview, XXVII:84–106
- Sandknop, E. M.: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182–193
- : Description of the larvae of the cusk eels *Ophidion scrippsae* and *Chilara taylora* (family Ophidiidae), XXIV:226–234
- : Development and distribution of larvae and pelagic juveniles of ocean whitefish, *Caulolatilus princeps*, in the CalCOFI survey region, XXVII:162–169
- Scannell, C. L.: The relative magnitude of the 1986 Pacific sardine spawning biomass off California, XXVIII:21–29
- Schmitt, P. D.: Prey size selectivity and feeding rate of larvae of the northern anchovy, *Engraulis mordax* Girard, XXVII:153–161
- Schultze, D. L.: California barracuda life history, fisheries, and management, XXIV:88–96
- Schwartzlose, R. A.: Large fluctuations in the sardine fishery in the Gulf of California: possible causes, XXVII:136–140
- Sciarrotta, T. C.: Abundance of southern California nearshore ichthyoplankton: 1978–1984, XXVII:53–64
- : Inshore soft substrata fishes in the Southern California Bight: an overview, XXVII:84–106
- Shelton, P.: Egg production estimates of anchovy biomass in the southern Benguela system, XXIX:137–157
- Shelton, P. A., A. J. Boyd, and M. J. Armstrong: The influence of large-scale environmental processes on neritic fish populations in the Benguela Current system, XXVI:72–92
- Siegel-Causey, D.: The temperate and tropical planktonic biotas of the Gulf of California, XXVII:228–266
- Singer, M. M.: Can we relate larval fish abundance to recruitment or population stability? A preliminary analysis of recruitment to a temperate rocky reef, XXVII:65–83
- : Inshore soft substrata fishes in the Southern California Bight: an overview, XXVII:84–106
- Smith, P. E.: Effects of sample size and contagion on estimating fish egg abundance, XXVIII:171–177
- : Geographical and seasonal patterns of larval fish species structure in the California Current area, 1975, XXIV:132–151
- : Ichthyoplankton and zooplankton abundance patterns in the California Current area, 1975, XXIV:109–131
- : An inverse egg production method for determining the relative magnitude of Pacific sardine spawning biomass off California, XXVI:130–138
- : Larval fish assemblages in the California Current region, 1954–1960, a period of dynamic environmental change, XXVIII:97–127
- : Recurrent groups of larval fish species in the California Current area, XXIV:152–164
- : The relative magnitude of the 1985 Pacific sardine spawning biomass off southern California, XXVII:25–31
- : The relative magnitude of the 1986 Pacific sardine spawning biomass off California, XXVIII:21–29
- Smith, P. E., and R. P. Hewitt: Anchovy egg dispersal and mortality as inferred from close-interval observations, XXVI:97–110

- Smith, P. E., and H. G. Moser: CalCOFI time series: an overview of fishes, XXIX:66-78
- Smith, S. E.: Trophic relationships of the shortbelly rockfish, *Sebastes jordani*, off central California, XXIX:129-136
- Stauffer, G. D.: Biology and fishery potential of jack mackerel, *Trachurus symmetricus*, XXIV:46-56
- : Calibration of northern anchovy spawning biomass estimates derived by egg production and larval census method, XXIV:11-15
- Stephens, J. S., Jr.: Inshore soft substrata fishes in the Southern California Bight: an overview, XXVII:84-106
- Stephens, J. S., Jr., G. A. Jordan, P. A. Morris, M. M. Singer, and G. E. McGowen: Can we relate larval fish abundance to recruitment or population stability? A preliminary analysis of recruitment to a temperate rocky reef, XXVII:65-83
- Stevens, E. G.: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182-193
- : Description of the larvae of the cusk eels *Ophidion scrippsae* and *Chilara taylori* (family Ophidiidae), XXIV:226-234
- : Development and distribution of larvae and pelagic juveniles of ocean whitefish, *Caulolatilus princeps*, in the CalCOFI survey region, XXVII:162-169
- Strange, E. M.: Drift bottle observations of the nearshore surface circulation off California, 1977-1983, XXV:68-73
- Stull, J. K., K. A. Dryden, and P. A. Gregory: A historical review of fisheries statistics and environmental and societal influences off the Palos Verdes Peninsula, California, XXVIII:135-154
- Sumida, B. Y.: Description of early stages of white seabass, *Atractoscion nobilis*, with notes on distribution, XXIV:182-193
- : Description of the larvae of the cusk eels *Ophidion scrippsae* and *Chilara taylori* (family Ophidiidae), XXIV:226-234
- : Development and distribution of larvae and pelagic juveniles of ocean whitefish, *Caulolatilus princeps*, in the CalCOFI survey region, XXVII:162-169
- Sumida, B. Y., and H. G. Moser: Food and feeding of bocaccio, *Sebastes paucispinis*, and comparison with Pacific hake, *Merluccius productus*, larvae in the California Current, XXV:112-118
- Sumida, B. Y., H. G. Moser, and E. H. Ahlstrom: Descriptions of larvae of California yellowtail, *Seriola lalandei*, and three other carangids from the eastern tropical Pacific: *Chloroscombrus orqueta*, *Caranx caballus*, and *Caranx sexfasciatus*, XXVI:139-159
- Sunada, J. S.: Spot prawn, *Pandalus platyceros*, and ridgeback prawn, *Sicyonia ingentis*, fisheries in the Santa Barbara Channel, XXV:100-104
- Theilacker, G. H., and A. S. Kimball: Comparative quality of rotifers and copepods as foods for larval fishes, XXV:81-86
- Thomas, D. H.: Year-to-year fluctuations of the California Countercurrent and effects on marine organisms, XXIV:165-181
- Trujillo-Ortíz, A.: Life cycle of the marine calanoid copepod, *Acartia californiensis* Trinast reared under laboratory conditions, XXVI:188-204
- Venrick, E. L., and T. L. Hayward: Determining chlorophyll on the 1984 CalCOFI surveys, XXV:74-79
- : Evaluation of some techniques for preserving nutrients in stored seawater samples, XXVI:160-168
- Vojkovich, M., and R. J. Reed: White seabass, *Atractoscion nobilis*, in California-Mexican waters: status of the fishery, XXIV:79-83
- Wolf, P., and P. E. Smith: An inverse egg production method for determining the relative magnitude of Pacific sardine spawning biomass off California, XXVI:130-138
- : The relative magnitude of the 1985 Pacific sardine spawning biomass off southern California, XXVII:25-31
- Wolf, P., P. E. Smith, and C. L. Scannell: The relative magnitude of the 1986 Pacific sardine spawning biomass off California, XXVIII:21-29
- Wosnitza-Mendo, C.: Peruvian hake fisheries from 1971 to 1982, XXVII:113-120

## SUBJECT INDEX

- Albacore  
age determination, XXIV:104  
feeding habits, XXVI:175–182  
fishery, XXIV:99–103  
1985, XXVII:13–14  
1986, XXVIII:18–19  
1987, XXIX:18  
migration patterns, XXIV:104  
oceanographic influences on, XXIV:104–106
- Anchovy, Benguela, *Engraulis capensis*  
distribution patterns, XXVI:72–92, XXIX:145–151  
spawning biomass, XXIX:137–157
- Anchovy, northern  
abundance, historical, XXIX:95–96  
egg dispersal vs mortality, XXVI:97–110  
and El Niño 1983, XXV:53–58  
fishery  
emergence of, XXIX:32–37  
1982, XXIV:6–7  
1983, XXV:8–9  
1984, XXVI:10  
1985, XXVII:8–9  
1986, XXVIII:12–13  
1987, XXIX:12–13  
larval feeding, ecology, and growth, XXVII:153–161  
larval feeding visual threshold, XXIV:245–254  
nearshore production and standing stock, XXIV:235–244  
spawning biomass  
calibration methods, XXIV:11–15  
1982–83, XXIV:16–27  
1983, XXV:16–27  
1984, XXVI:17–25  
1985, XXVII:16–24  
spawning habitat  
entrainment by eddy, 1985, XXVII:144–152  
satellite remote sensing of, XXIV:202–209
- Anchovy, Peruvian, *Engraulis ringens*  
sex ratio, XXV:50–51  
spawning frequency, XXV:43–50
- Barracuda, *Sphyræna argentea*  
life history and fishery, XXIV:88–92
- Bass  
barred sand, *Paralabrax nebulifer*  
batch fecundity and ovary subsampling, XXVIII:168–170  
feeding habits, XXV:105–111  
kelp, *Paralabrax clathratus*  
batch fecundity and ovary subsampling, XXVIII:168–170
- Benguela anchovy. *See* Anchovy, Benguela
- Benguela Current system  
anomalous warm events in, XXVI:51–63  
and neritic fish, XXVI:72–92
- Bight, Southern California  
*Euphausia pacifica* in, 1977–78, XXV:87–99  
inshore soft substrata fishes, XXVII:84–104  
vertical distribution of biomass and species, XXIV:210–215
- Billfish fishery, XXIV:70–78. *See also* Swordfish
- Bluefin tuna, *Thunnus thynnus*  
fisheries, XXIV:97–98
- Blue rockfish, *Sebastes mystinus*  
growth, XXIV:216–225
- Bocaccio, *Sebastes paucispinis*  
feeding habits, XXV:112–118
- Bonito, *Sarda chiliensis*  
feeding habits, XXVI:175–182
- CalCOFI  
history, XXIX:27–65  
time series, overview of fishes, XXIX:66–77
- California halibut. *See* Halibut
- Canal de Ballenas  
near-surface currents in, XXV:35–42
- Carangid larvae, XXVI:139–159  
*Caranx caballus*, XXVI:149–152, 157–159  
*Caranx sexfasciatus*, XXVI:152–159  
*Chloroscombrus orqueta*, XXVI:143–149, 157–159  
*Seriola lalandei*, XXVI:139–142, 157–159
- Chlorophyll determination, XXV:74–79
- Clark, F. N. In memoriam, XXVIII:5
- Continental shelf, southern California  
physical-chemical characteristics and zooplankton biomass, XXVII:36–52
- Copepods  
*Acartia californiensis* life cycle, XXVII:188–204  
community structure and biomass during El Niño, XXIX:122–128  
in eastern equatorial Pacific, XXVII:170–187  
in eastern tropical Pacific, XXVII:205–227  
as food for larval fishes, XXV:80–85  
in Gulf of California, XXVII:242–246
- Countercurrent, California  
effects on marine organisms, XXIV:178–180  
year-to-year fluctuations, XXIV:165–178
- Crab  
Dungeness, *Cancer magister*, fishery  
1982, XXIV:9–10  
1983, XXV:12  
1984, XXVI:13–14  
1985, XXVII:11–12  
1986, XXVIII:15–16  
1987, XXIX:16  
pelagic red and California Countercurrent, XXIV:178–179
- Current systems. *See* Benguela Current system; Eastern boundary current systems
- Cusk eels, *Chilara taylori* and *Ophidion scrippsae*  
larvae, XXIV:226–234
- Data system, CalCOFI, XXIX:38–40
- Demersal fishes  
catch  
diel variation in, XXV:119–134  
1982, XXIV:9  
1983, XXV:11–12  
1984, XXVI:12–13  
1985, XXVII:11  
1986, XXVIII:15  
1987, XXIX:15–16  
species composition and abundance, XXV:119–134, XXVIII:155–167
- Dungeness crab. *See* Crab
- Eastern boundary current systems, XXVI:30–40, 51–63, 72–91
- Eggs, fish  
anchovy, dispersal of, XXVI:97–110  
cusk eels, XXIV:226–234  
estimating abundance of, XXVIII:171–177  
sampling with CalVET net, XXVIII:178–182  
white seabass, XXIV:183

- El Niño  
  copepods during, XXIX:122–128  
  and larval fish abundance in King Harbor, Calif., XXVII:65–83  
  in nearshore zone, XXVII:48–49  
  1957–58, physical oceanography after, XXIX:56–63  
  and productivity anomalies XXVI:65–71
- Engraulis*. See Anchovy
- Euphausia pacifica*  
  larval morphogenesis, 1977–78, XXV:87–99  
  See also Plankton
- Fisheries  
  CalCOFI time series, XXIX:66–77  
  oceanographic approach to research, XXIX:27–41  
  review  
    1982, XXIV:6–10  
    1983, XXV:7–15  
    1984, XXVI:9–16  
    1985, XXVII:7–15  
    1986, XXVIII:11–20  
    1987, XXIX:11–20  
  See also individual fishes
- Fleminger, A. In memoriam, XXIX:5–6
- Frame trawl, XXVII:267–278
- Fronts, observed with infrared scanner, XXV:59–67
- Fur seals  
  food consumption, XXV:135–145  
  population estimates, XXV:137, 138–140
- Groundfish. See Demersal fishes
- Gulf of California  
  copepods in, XXIX:122–127  
  fisheries  
    history, XXVIII:32–36  
    sardine, XXVII:136–140, XXVIII:128–134  
  zooplankton biomass in, XXIX:122–127
- Hake, Pacific, *Merluccius productus*  
  abundance, historical, XXIX:98–99  
  feeding habits, XXV:112–118  
  recruitment forecasting, XXVII:109–120  
  spawning biomass, XXIX:81–90
- Hake, Peruvian, *Merluccius gayi peruanus*  
  fisheries, 1971–82, XXVII:113–120
- Halibut, California, *Paralichthys californicus*  
  catch  
    1986, XXVIII:17  
    1987, XXIX:17  
  distribution, XXIV:194, 200–201  
  feeding habits, XXIV:196–204  
  size limit, effect on fisheries, XXIX:159–166
- Herring, Pacific, *Clupea harengus*  
  catch  
    1982, XXIV:8–9  
    1983, XXV:11  
    1984, XXVI:12  
    1985, XXVII:10–11  
    1986, XXVIII:14–15  
    1987, XXIX:15  
  See also Thread herring
- Ichthyoplankton. See Plankton
- Infrared scanner  
  and coastal fronts, XXV:59–67
- In memoriam  
  Clark, F. N., XXVIII:5  
  Fleminger, A., XXIX:5–6  
  Johnson, M. W., XXVI:5–6  
  Lasker, R., XXIX:7–8  
  Roedel, P. M., XXVIII:6  
  Sargent, M., XXVIII:7
- Jack mackerel. See Mackerel, jack
- Johnson, M. W. In memoriam, XXVI:5–6
- Kelp bass, *Paralabrax clathratus*  
  batch fecundity, XXVIII:168–170
- Larval fish  
  abundance and recruitment or population stability, XXVII:65–83  
  King Harbor, Calif., populations (1974–85), XXVII:67–70  
  assemblages in California Current area, 1954–60, XXVIII:97–127  
  foods, comparative quality of, XXV:80–86  
  geographical and seasonal patterns, 1975, XXIV:132–151  
  recurrent groups in California Current area, XXIV:152–164, XXVIII:101–120  
  *Sebastes*, abundance of four species, XXVII:121–135  
  time series, community and population processes, XXIX:66–77  
  See also Plankton
- Lasker, R. In memoriam, XXIX:7–8.
- Lobster, California spiny, *Panulirus interruptus*  
  catch  
    1984, XXVI:15–16  
    1985, XXVII:12–13  
    1986, XXVIII:17–18  
    1987, XXIX:17–18
- Mackerel, jack, *Trachurus symmetricus*  
  abundance, historical, XXIX:98  
  biology, XXIV:46–48  
  catch  
    1982, XXIV:7  
    1983, XXV:9  
    1984, XXVI:10–11  
    1985, XXVII:9  
    1986, XXVIII:13  
    1987, XXIX:13  
  fishery potential, XXIV:46–56  
  growth in captivity, XXV:146–150
- Mackerel, Pacific, *Scomber japonicus*  
  abundance, XXVI:119–129, XXIX:96–97  
  catch  
    1982, XXIV:7–8  
    1983, XXV:9–10  
    1984, XXVI:11–12  
    1985, XXVII:9–10  
    1986, XXVIII:13–14  
    1987, XXIX:13–14  
  fishery and management history, XXIV:35–45  
  spawning biomass, 1929–69, XXIX:81–90
- Marlin, striped, fishery, XXIV:70, 75–78
- Memorials. See In memoriam
- Merluccius. See Hake
- Neritic fish  
  in Benguela Current system, XXVI:72–92
- Northern anchovy. See Anchovy, northern
- Ocean whitefish. See Whitefish
- Opisthonema*. See Thread herring
- Pacific hake. See Hake, Pacific
- Pacific herring. See Herring, Pacific
- Pacific sardine. See Sardine
- Pacific saury. See Saury
- Pacific whiting. See Hake, Pacific
- Palos Verdes Peninsula fisheries, XXVIII:135–154

- Paralabrax*. See Bass  
Peruvian hake. See Hake, Peruvian  
Physical oceanography  
California Countercurrent fluctuations, XXIV:165–180  
coastal fronts observed with infrared scanner, XXV:59–67  
dynamics affecting eastern tropical Atlantic and Pacific, XXVI:41–50  
entrainment of anchovy spawning habitat, XXVII:144–152  
mesoscale cycles in Gulf of California, XXVIII:128–134  
near-surface currents in Canal de Ballenas, XXV:35–42  
1947–87, XXIX:42–65  
nutrient concentration in upwelling regions, XXVI:93–96  
physical-chemical characteristics on southern California continental shelf, XXVII:36–52  
surface circulation off California, 1977–83, XXV:68–73  
warm events in eastern boundary currents, XXVI:51–64  
Plankton  
in Gulf of California, XXVII:228–266  
ichthyoplankton  
larval growth off Peru, XXVI:113–118  
nearshore abundance, XXVII:53–64  
seasonal changes in California Current, 1975, XXIV:109–131  
zooplankton  
seasonal variability in California Current, XXIV:109–131, XXVIII:72–80  
See also Larval fish  
Recreational fishery  
1983, XXV:13–15  
1985, XXVII:14  
1986, XXVIII:19  
1987, XXIX:19–20  
Recruitment problem, comparative studies, XXVI:30–40  
Ridgeback prawn, *Sicyonia ingentis*  
review of 1987 fishery, XXIX:18–19  
in Santa Barbara channel, XXV:100–104  
Rockfish. See *Sebastes*  
Roedel, P. M. In memoriam, XXVIII:6  
Rotifers  
as food for larval fishes, XXV:80–86  
Sample size and contagion  
effects on estimating egg abundance, XXVIII:171–177  
San Pedro fishery, XXIV:255–258  
Sardine, Pacific, *Sardinops sagax*  
abundance, historical, XXIX:96  
catch  
1982, XXIV:6  
1983, XXV:7–8  
1984, XXVI:9–10  
1985, XXVII:7–8  
1986, XXVIII:11–12  
1987, XXIX:11–12  
in Gulf of California, XXVIII:128–134, XXIX:102–109  
larval growth, XXVI:113–118  
research history, XXIX:28–37  
satellite imagery, XXIX:103  
spawning biomass  
determining, XXVI:130–138  
1985, XXVII:25–31  
1986, XXVIII:21–26  
Sardines  
Gulf of California fishery, XXVII:136–140, XXVIII:128–134  
Sargent, M. In memoriam, XXVIII:7  
Saury, Pacific, *Cololabis saira*  
abundance, historical, XXIX:99  
Seabass, white, *Atractoscion nobilis*  
distribution, XXIV:192  
early stages, XXIV:182–192  
fishery, XXIV:79–83  
Seals. See Fur seals  
Seawater  
preserving nutrients in, XXVI:160–168  
*Sebastes*  
*jordani*, trophic relationships, XXIX:129–136  
larval abundance of four species, XXVII:121–125  
*mystinus*, growth, XXIV:216–225  
*paucispinis*, feeding habits, XXV:112–118  
Sharks, pelagic  
biology and fisheries of common thresher, bonito, and blue, XXIV:57–69  
catch  
1982, XXIV:10  
1983, XXV:13  
1984, XXVI:14–15  
1985, XXVII:12  
1986, XXVIII:16–17  
1987, XXIX:17  
Shortbelly rockfish, *Sebastes jordani*  
diet and prey distribution, XXIX:132–135  
trophic relationships, XXIX:129–136  
Shrimp fisheries  
in Mexico, XXVIII:43–52  
review  
1983, XXV:12–13  
1984, XXVI:14  
1985, XXVII:12  
1986, XXVIII:16  
1987, XXIX:16–17  
Skipjack  
feeding habits, XXVI:175–182  
See also Tuna  
Southern California Bight. See Bight, Southern California  
Spiny lobster. See Lobster  
Spot prawn, *Pandalus platyceros*, fisheries  
1987, XXIX:18–19  
in Santa Barbara Channel, XXV:100–104  
Squid, *Loligo opalescens*, fishery  
1982, XXIV:8  
1983, XXV:10–11  
1984, XXVI:12  
1985, XXVII:10  
1986, XXVIII:14  
1987, XXIX:14–15  
Striped marlin, *Tetrapturus audax*, fishery, XXIV:70, 75–78  
Surface circulation off California, 1977–83, XXV:68–73  
Swordfish, *Xiphias gladius*, fishery  
to 1981, XXIV:70–75, 78  
1982, XXIV:10  
1983, XXV:13  
1984, XXVI:14–15  
1985, XXVII:12  
1986, XXVIII:16–17  
1987, XXIX:17  
Thread herring, *Opisthonema*  
fishery fluctuations, XXVII:136–140  
genetic and morphological divergence of three species, XXIX:110–121  
Time series, CalCOFI  
overview of fishes, XXIX:66–77  
Trawl for juvenile fishes, XXVII:267–278



Tuna fisheries

eastern Pacific Ocean, XXIV:97–98, XXVIII:37–38

global overview, XXVIII:37

Mexican, XXVIII:39–42

*See also common names*

Upwelling regions

nutrient concentration in, XXVI:93–96

Whitefish, ocean, *Caulolatilus princeps*

larvae and juveniles, XXVII:162–169

White seabass. *See* Seabass, white

Whiting. *See* Hake

Yellowtail, *Seriola lalandei*

catch analysis and life history, XXIV:84–87

larvae, XXVI:139–142, 157–159